

Disparities in Testing for Deficient Mismatch Repair Genes Among Medicare Beneficiaries With Colon Cancer in the United States: a claims-based analysis.

Amanda Speller, Eunice Chang, Caitlin Sheetz, Michael S. Broder
ADVI Health, Washington, DC


ISPOR 2025

May 13 – 16, 2025
Montreal, QC, Canada
Poster Code: HPR35
Abstract Number: 5091

Background

- Guidelines recommend deficient mismatch repair/ microsatellite instability (dMMR/MSI) genomic testing for all patients with colon cancer.¹
- dMMR/MSI is present in 5% - 20% of colon cancers.²
- Patients with dMMR/MSI positive tumors benefit from immunotherapies that help the immune system attack cancer cells.³
- The benefits of innovative immunotherapies tend to reach socially disadvantaged groups last.⁴

State-level variation in dMMR/MSI genomic testing
30 days prior to or 90 days after a colon cancer diagnosis



Why is it important to know whether patients are tested for dMMR/MSI?


- Inadequate genomic testing may worsen disparities in immunotherapy use and, in turn, reduce survival.
- Identifying patient groups who are less likely to receive dMMR/MSI genomic testing is a critical step towards increasing guideline-based screening and treatment.

Almost half (43.4%) of patients diagnosed with colon cancer *did not* receive guideline-based genomic testing. Patients who received guideline-based care were more likely to use life-saving immunotherapies (4.9% versus 1.8%, $p < 0.001$).

Data and methods

- Used the 100% Medicare Fee-For-Service Research Identifiable Files (RIFs).
- Identified patients ($n=20,809$) who were diagnosed with colon cancer in 2022.
 - In this cohort, 18.6% received a low-income subsidy and 84.1% were Non-Hispanic White.
- Measured receipt of genomic testing and utilization of immunotherapy.
- Tested differences in the likelihood of dMMR/MSI genomic testing by patient sociodemographic characteristics.

Relative risk of dMMR/MSI genomic testing by patient sociodemographic characteristics



What does this study mean for patients with colon cancer?

- This study identified patients at risk for poor outcomes due to low quality care.
- Patients who received a low-income subsidy or who were non-White were less likely to receive dMMR/MSI testing. Reducing testing barriers for these groups may also reduce survival disparities.
- Geographic disparities in dMMR/MSI testing suggests that non-clinical factors influence testing rates. State-level initiatives may help increase uptake.

References

- Hechtman JF, et al. Universal screening for microsatellite instability in colorectal cancer in the clinical genomics era: new recommendations, methods, and considerations. *Fam Cancer*. 2017 Oct;16:525-9.
- Taieb J, et al. Deficient mismatch repair/microsatellite unstable colorectal cancer: Diagnosis, prognosis and treatment. *Eur J Cancer*. 2022 Nov;175:136-57.
- Andre T, et al. Nivolumab plus Ipilimumab in Microsatellite-Instability–High Metastatic Colorectal Cancer. *N Engl J Med*. 2024 Nov;391(21):2014-26.
- Carroll CE, et al. Adoption of innovative therapies across oncology practices—evidence from immunotherapy. *JAMA Oncol*. 2023 Mar;9(3):324-33.

Amanda Speller

Associate Director | ADVI Health

Email | amanda.speller@advi.com

